



**ADVANCED SUBSIDIARY GCE  
PHYSICS A**  
Forces and Motion

**2821**

**Tuesday 13 January 2009  
Afternoon**

**Duration: 1 hour**

Candidates answer on the question paper

**OCR Supplied Materials:**  
None

**Other Materials Required:**

- Electronic calculator
- Ruler
- Protractor



Candidate Forename	Cowen	Candidate Surname	Physics
--------------------	-------	-------------------	---------

Centre Number				Candidate Number			
---------------	--	--	--	------------------	--	--	--

**INSTRUCTIONS TO CANDIDATES**

- Write your name clearly in capital letters, your Centre Number and Candidate Number in the boxes above.
- Use black ink. Pencil may be used for graphs and diagrams only.
- Read each question carefully and make sure that you know what you have to do before starting your answer.
- Answer all the questions.
- Do not write in the bar codes.
- Write your answer to each question in the space provided, however additional paper may be used if necessary.

**INFORMATION FOR CANDIDATES**

- The number of marks is given in brackets [ ] at the end of each question or part question.
- The total number of marks for this paper is 60.
- You will be awarded marks for the quality of written communication where this is indicated in the question.
- You may use an electronic calculator.
- You are advised to show all the steps in any calculations.
- This document consists of 16 pages. Any blank pages are indicated.

FOR EXAMINER'S USE		
Qu.	Max.	Mark
1	16	
2	8	
3	10	
4	6	
5	10	
6	10	
<b>TOTAL</b>	<b>60</b>	

**Data**

- speed of light in free space,  $c = 3.00 \times 10^8 \text{ m s}^{-1}$
- permeability of free space,  $\mu_0 = 4\pi \times 10^{-7} \text{ H m}^{-1}$
- permittivity of free space,  $\epsilon_0 = 8.85 \times 10^{-12} \text{ F m}^{-1}$
- elementary charge,  $e = 1.60 \times 10^{-19} \text{ C}$
- the Planck constant,  $h = 6.63 \times 10^{-34} \text{ J s}$
- unified atomic mass constant,  $u = 1.66 \times 10^{-27} \text{ kg}$
- rest mass of electron,  $m_e = 9.11 \times 10^{-31} \text{ kg}$
- rest mass of proton,  $m_p = 1.67 \times 10^{-27} \text{ kg}$
- molar gas constant,  $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
- the Avogadro constant,  $N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$
- gravitational constant,  $G = 6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$
- acceleration of free fall,  $g = 9.81 \text{ m s}^{-2}$

## Formulae

uniformly accelerated motion,

$$s = ut + \frac{1}{2} at^2$$

$$v^2 = u^2 + 2as$$

refractive index,

$$n = \frac{1}{\sin C}$$

capacitors in series,

$$\frac{1}{C} = \frac{1}{C_1} + \frac{1}{C_2} + \dots$$

capacitors in parallel,

$$C = C_1 + C_2 + \dots$$

capacitor discharge,

$$x = x_0 e^{-t/CR}$$

pressure of an ideal gas,

$$p = \frac{1}{3} \frac{Nm}{V} \langle c^2 \rangle$$

radioactive decay,

$$x = x_0 e^{-\lambda t}$$

$$t_{1/2} = \frac{0.693}{\lambda}$$

critical density of matter in the Universe,

$$\rho_0 = \frac{3H_0^2}{8\pi G}$$

relativity factor,

$$= \sqrt{1 - \frac{v^2}{c^2}}$$

current,

$$I = nAve$$

nuclear radius,

$$r = r_0 A^{1/3}$$

sound intensity level,

$$= 10 \lg \left( \frac{I}{I_0} \right)$$

Answer all the questions.

- 1 A projectile is fired, from ground level, into the air at a velocity of  $50 \text{ m s}^{-1}$  and  $53^\circ$  to the horizontal. Fig. 1.1 shows the variation of the vertical component of the velocity  $v_v$  against time,  $t$ , until the projectile reaches its maximum height at 4.1 s.




Fig. 1.1

Fig. 1.2 shows the variation of the horizontal component of the velocity  $v_h$  against time,  $t$ , for the same part of the journey.




Fig. 1.2

(a) (i) Use Figs. 1.1 and 1.2 to describe the motion of the projectile

1 in the vertical direction

The projectile decelerates at a constant rate [1]

2 in the horizontal direction.

Constant velocity [1]

(ii) Use Figs. 1.1 and 1.2 to calculate at the maximum height

1 the vertical displacement

Area under graph:  $\frac{40 \times 4.1}{2} = 82$

vertical displacement = 82 m [2]

2 the horizontal displacement.

Area under graph:  $\frac{30 \times 4.1}{2} = 61.5$  ~~123~~

horizontal displacement = 61.5 m [1]

(b) Describe the difference between the distance travelled by the projectile and the displacement of the projectile.


The projectile's ~~has~~ distance is the arc that it has travelled along, the displacement is the shorter distance length in a straight line from start to end. [2]

(c) (i) Use Figs. 1.1 and 1.2 to determine  $v_v$  and  $v_h$  after 2.0s.

$v_v = 20 \text{ ms}^{-1}$

$v_h = 30 \text{ ms}^{-1}$  [1]

(ii) Hence or otherwise calculate the velocity of the projectile after 2.0s.



$$20^2 + 30^2 = 36.1$$

$$\tan \theta = \frac{20}{30}$$

$$\theta = \tan^{-1}\left(\frac{20}{30}\right) = 33.7^\circ$$

magnitude of the velocity = 36.1  $\text{ms}^{-1}$

angle of velocity to the horizontal = 33.7  $^\circ$  [4]

(iii) State the magnitude of the velocity of the projectile after 4.1s.

velocity = 30  $\text{ms}^{-1}$  [1]

(d) Continue the graphs in Figs. 1.1 and 1.2 to show how the velocity of the projectile varies up to 6.0s. [3]

[Total: 16]

PLEASE DO NOT WRITE ON THIS PAGE

2 Fig. 2.1 shows apparatus used to determine the acceleration due to gravity.




Fig. 2.1

A clock starts when a ball is released from an electromagnet and stops when the ball opens a trapdoor. The clock then displays the time  $t$  for the ball to fall a distance  $s$ . The trapdoor is moved to change the value of  $s$ . Fig. 2.2 shows the variation of  $t^2$  against distance  $s$ .




Fig. 2.2

- (a) Use an equation of motion to explain why the graph in Fig. 2.2 is a straight line.

$$y = mx + c \quad s = ut + \frac{1}{2}at^2$$

$$t^2 = \frac{2as}{a} \quad \text{Gradient} = \frac{2a}{a}$$

[2]

- (b) Use Fig. 2.2 to calculate

- (i) the gradient of the graph

$$m = \frac{0.1}{0.5} = 0.20$$

gradient =  $0.20$  [2]

- (ii) the acceleration of the ball.

$$0.2 = \frac{2}{a} \quad a = \frac{2}{0.2} = 10.0$$

acceleration =  $10.0$   $\text{ms}^{-2}$  [2]

- (c) When a student performs this experiment he obtains a straight line graph which does not pass through the origin. Suggest **two** reasons why.

Air resistance.

Time delay releasing ball.

[2]

[Total: 8]

- 3 (a) Define

- (i) the newton

The force required to accelerate 1kg by  $1\text{ms}^{-2}$  [1]

- (ii) work done by a force.

Work done by a force is equal to the force multiplied by the distance moved in the direction of the force. [1]

- (b) A car of mass  $m$  is travelling at a speed  $u$ . A constant braking force  $F$  is applied until the car comes to rest. The car travels a distance  $s$  after the force is applied.

- (i) Write equations for

- 1 the work done by the force  $F$

$$W = Fs$$

- 2 the relationship between the force  $F$  and acceleration  $a$ .

$$F = ma$$

[1]

- (ii) Use the equations in (i) to show that the work done bringing the car to rest equals the loss in kinetic energy of the car.

$$E_k = \frac{1}{2}mu^2 \quad W = mas$$

$$v^2 = u^2 + 2as$$

$$v = 0$$

$$u^2 = -2as$$

$$as = -\frac{u^2}{2}$$

$$W = -m \frac{u^2}{2} = -\frac{1}{2}mu^2 = E_k$$

[3]

(iii) Hence describe the relationship between the braking distance and the velocity of a car.

*Braking distance is proportional to square of velocity.*

..... [1]

(c) (i) A car of mass 1550 kg has a braking distance of 25.0 m when travelling at 13.3 ms<sup>-1</sup>. Calculate the average braking force acting on the car.

*s = 25, u = 13.3, v = 0, a = ?*

*F = ma = 1550 × 3.5 = 5500*

*v<sup>2</sup> = u<sup>2</sup> + 2as*

*a =  $\frac{v^2 - u^2}{2s} = \frac{-13.3^2}{2 \times 25} = -3.5 \text{ ms}^{-2}$*

*force = 5500 N [2]*

(ii) Deduce the braking distance when the car is travelling at 26.6 ms<sup>-1</sup> and the same braking force is applied.

*25 = k 13.3<sup>2</sup>*

*k =  $\frac{25}{13.3^2} = 0.141$  distance = 100 m [1]*

*26.6<sup>2</sup> × 0.141 = 100*

[Total: 10]

4 (a) (i) Define pressure.

*Force per unit area*

..... [1]

(ii) Define density.

*Mass per unit volume*

..... [1]

(b) Fig. 4.1 shows the outline of a swimming pool that is on the top floor of a building.




Fig. 4.1

The total volume of concrete used is 27 m<sup>3</sup>. The density of the concrete is 2600 kg m<sup>-3</sup>.

(i) Calculate the mass of concrete.

*2600 × 27 = 70200*

mass = *70200* kg [1]

- (ii) The pool contains water to a constant depth of 1.5 m. The internal dimensions of the pool are 8.0 m by 12.0 m. The density of water is  $1000 \text{ kg m}^{-3}$ . Calculate the pressure the water exerts on the base of the pool.

$$V = 1.5 \times 8 \times 12 = 144 \text{ m}^3$$

$$m = 144 \times 1000 = 144000 \text{ kg}$$

$$p = \frac{F}{A} = \frac{mg}{A} = \frac{144000 \times 9.81}{8 \times 12} = 14.7 \text{ kPa}$$

pressure = 14.7 kPa [2]

- (iii) Suggest with a reason whether the pressure on the floor under the concrete walls or under the water is greater.

The concrete because it is more dense and has higher walls.

[1]

[Total: 6]

- 5 (a) Define the centre of gravity of a body.

The point about which the weight of an object can be assumed to act. [1]

Fig. 5.1 shows a boat being lifted vertically. It is attached to two ropes labelled 1 and 2. A motor lifts the boat vertically.




Fig. 5.1

- (b) When the boat is rising at a constant speed the tensions in ropes 1 and 2 are  $T_1$  and  $T_2$  respectively. The weight of the boat is 22 kN. The weight acts 1.20 m and 0.40 m from the lines of action of the tensions  $T_1$  and  $T_2$  respectively.

- (i) Explain why the rising boat can be described as being in equilibrium.

Travelling at constant speed/velocity.

[1]

- (ii) Calculate the moment of the weight of the boat about the point P.

$C = G: 1.6 \text{ m}$

$$\text{moment} = 0.4 \times 22000 = 8800 \text{ Nm}$$

moment = 8800 Nm unit [2]

- (iii) Calculate the tension  $T_1$ .

$C = G: 1.6 \text{ m}$

$$T_1 = \frac{8800}{1.6} = 5500 \text{ N}$$

$T_1 = 5500 \text{ N}$  [2]

- (iv) Calculate the tension
- $T_2$
- .

$$5500 + T_2 = 22000$$

$$T_2 = 22000 - 5500 = 16500$$

$$T_2 = 16500 \text{ N [1]}$$

- (v) Calculate the minimum power required by the motor to lift the boat at a constant speed of
- $0.015 \text{ ms}^{-1}$
- .

$$P = Fv = 22000 \times 0.015 \\ = 330 \text{ W}$$

$$\text{power} = 330 \text{ W [2]}$$

- (c) Describe and explain one situation during the lifting of the boat when it will not be in equilibrium.

When it is accelerating at start  
~~to start~~

[1]

[Total: 10]

6 In this question, two marks are available for the quality of written communication.

- (a) Use the following terms to explain the behaviour of a spring when a tensile force is applied:

extension      Hooke's law      elastic limit      spring constant

Extension is the increase in the spring length when a force is applied.

Hooke's law: Force is proportional to extension, as long as elastic limit is not exceeded.

Elastic limit is the maximum force before the spring suffers permanent deformation.

[3]

THIS QUESTION CONTINUES ON PAGE 16

$$\text{Force constant} = k = \frac{\text{force}}{\text{extension}} \quad \text{[Turn over]}$$

- (b) Describe how the following terms may be used to compare materials:

stiff      strong      brittle      ductile

In your answer you should include the quantities stress, strain and Young modulus and the terms elastic and plastic.

Stiff material has a high Young modulus.

Strong material can take a high stress before breaking.

Brittle materials have no plastic deformation.

Ductile materials have large plastic region.

$$\text{Young modulus} = \frac{\text{stress}}{\text{strain}}$$

[5]

Quality of Written Communication [2]

[Total: 10]

END OF QUESTION PAPER

**OCR**   
RECOGNISING ACHIEVEMENT

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (OCR) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

OCR is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.